

Complex Numbers

Paul Renteln

Department of Physics
California State University
San Bernardino, CA 92407

May, 2009

1 The Basic Representation

We define the imaginary quantity i to be

$$i = \sqrt{-1}. \quad (1.1)$$

Any complex number can be written in the *standard form*

$$z = a + ib, \quad (1.2)$$

where a and b are real numbers. The *real part* of z , denoted $\operatorname{Re} z$, is the *real* number a . The *imaginary part* of z , denoted $\operatorname{Im} z$, is the *real* number b . The *complex conjugate* of z , written z^* (or sometimes \bar{z}), is the complex number

obtained from z by flipping the sign of the imaginary part:

$$z = a + ib \quad \Rightarrow \quad z^* = a - ib. \quad (1.3)$$

Complex numbers are added, subtracted, multiplied, and divided just like real numbers. So

$$(a + ib) + (c + id) = (a + c) + i(b + d), \quad (1.4)$$

and

$$(a + ib)(c + id) = (ac - bd) + i(ad + bc). \quad (1.5)$$

The ratio of two complex numbers can be put into standard form by multiplying the numerator and denominator by the complex conjugate of the denominator:

$$\begin{aligned} \frac{a + ib}{c + id} &= \frac{a + ib}{c + id} \left(\frac{c - id}{c - id} \right) \\ &= \frac{(ac + bd) + i(ad - bc)}{c^2 + d^2} \\ &= \left(\frac{ac + bd}{c^2 + d^2} \right) + i \left(\frac{ad - bc}{c^2 + d^2} \right). \end{aligned} \quad (1.6)$$

2 The Modulus

Note that the product of a complex number and its complex conjugate is always a real number:

$$z = a + ib \quad \Rightarrow \quad zz^* = a^2 + b^2. \quad (2.1)$$




Figure 1: The complex plane

The *modulus* of z , written $|z|$, is defined to be the *positive root* of zz^* :

$$|z| = (zz^*)^{1/2}. \quad (2.2)$$

A short calculation shows that the modulus is multiplicative, meaning that, for any two complex numbers z_1 and z_2 ,

$$|z_1 z_2| = |z_1| |z_2|. \quad (2.3)$$

Similarly,

$$\left| \frac{z_1}{z_2} \right| = \frac{|z_1|}{|z_2|}. \quad (2.4)$$

3 A Geometric Interpretation

There is a one-to-one correspondence between complex numbers and points of the plane, given by

$$a + ib \quad \longleftrightarrow \quad (a, b), \quad (3.1)$$

and illustrated in Figure 1.

This allows us to specify a complex number by its polar coordinates, usually denoted (r, θ) . From elementary trigonometry it is clear that

$$a = r \cos \theta \qquad r = (a^2 + b^2)^{1/2} = |z| \qquad (3.2)$$

$$b = r \sin \theta \qquad \theta = \tan^{-1} \left(\frac{b}{a} \right). \qquad (3.3)$$

r (which is just the modulus of z) is sometimes called the *length* of z , and θ is called the *argument* (or *angle*) of z , and is denoted $\text{Arg } z$.

4 The Complex Exponential Representation

Complex numbers admit another representation in terms of complex exponentials. To motivate this, recall that, near $x = 0$, a differentiable function can be approximated by its MacLaurin-Taylor expansion

$$f(x) = f(0) + f'(0)x + f''(0)\frac{x^2}{2!} + f'''(0)\frac{x^3}{3!} + \dots \qquad (4.1)$$

Applying this formula enables us to write

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots \qquad (4.2)$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} + \dots \qquad (4.3)$$

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} + \dots \qquad (4.4)$$

Substituting $x = i\theta$ in the expansion of e^x and collecting real and imaginary parts gives

$$\begin{aligned} e^{i\theta} &= 1 + i\theta + \frac{(i\theta)^2}{2!} + \frac{(i\theta)^3}{3!} + \dots \\ &= \left(1 - \frac{\theta^2}{2!} + \frac{\theta^4}{4!} + \dots\right) + i \left(\theta - \frac{\theta^3}{3!} + \frac{\theta^5}{5!} + \dots\right), \end{aligned}$$

whereupon we obtain *Euler's formula*:

$$e^{i\theta} = \cos \theta + i \sin \theta. \quad (4.5)$$

Among other things, Euler's formula yields the classic equality

$$e^{i\pi} = -1.$$

Using (3.2) and (3.3) we can write any complex number as

$$z = a + ib = r \cos \theta + ir \sin \theta, \quad (4.6)$$

so comparing with (4.5) we see that any complex number can be written in *complex exponential form*

$$z = re^{i\theta}. \quad (4.7)$$

As a consequence of Euler's formula we get the very useful equations:

$$\cos \theta = \frac{1}{2} (e^{i\theta} + e^{-i\theta}) \quad (4.8)$$

$$\sin \theta = \frac{1}{2i} (e^{i\theta} - e^{-i\theta}). \quad (4.9)$$

The complex exponential representation is ideally suited to multiplication

and division of complex numbers because of the corresponding properties of the exponential. In particular, for any two complex numbers

$$z_1 = r_1 e^{i\theta_1} \quad \text{and} \quad z_2 = r_2 e^{i\theta_2}, \quad (4.10)$$

we have

$$z_1 z_2 = r_1 r_2 e^{i(\theta_1 + \theta_2)} \quad \text{and} \quad \frac{z_1}{z_2} = \frac{r_1}{r_2} e^{i(\theta_1 - \theta_2)}. \quad (4.11)$$

Geometrically, the product of z_1 and z_2 has length equal to the product of the lengths of z_1 and z_2 and argument equal to the sum of the arguments of z_1 and z_2 .

5 Roots of Unity

According to the Fundamental Theorem of Algebra, an n^{th} order polynomial has n complex roots. This means that the equation

$$z^n = 1 \quad (5.1)$$

has n solutions, which are called n^{th} *roots of unity*. To find them, we use Euler's formula (4.5) to observe that, for any integer k ,

$$e^{2\pi ki} = 1. \quad (5.2)$$

Thus (5.1) can be solved by writing

$$z = (1)^{1/n} = (e^{2\pi ki})^{1/n} = e^{2\pi(k/n)i}. \quad (5.3)$$




Figure 2: The three roots of unity

Observe that we get distinct solutions only for the values $k = 0, 1, 2, \dots, n-1$, after which the solutions repeat. Equivalently, we get distinct solutions only for the integers modulo n .¹ Hence, the n^{th} roots of unity are

$$1, e^{2\pi i/n}, e^{4\pi i/n}, e^{6\pi i/n}, \dots, e^{2(n-1)\pi i/n}. \quad (5.4)$$

Example 1 The cube roots of unity are

$$1, e^{2\pi i/3}, \text{ and } e^{4\pi i/3}.$$

Equivalently, they are

$$1, -\frac{1}{2} + i\frac{\sqrt{3}}{2}, \text{ and } -\frac{1}{2} - i\frac{\sqrt{3}}{2}. \quad (5.5)$$

(See Figure 2.) Note that they are evenly spaced around the unit circle.

¹Recall that, if a and b are integers and n is a natural number, we say a is congruent to b modulo n , written $a \equiv b \pmod{n}$, if $a - b$ is evenly divisible by n .

Example 2 The cube roots of $2i$ can be obtained by rewriting $2i$ in complex exponential representation:

$$(2i)^{1/3} = \left(2e^{i\pi/2+2\pi ki}\right)^{1/3} = 2^{1/3}e^{i\pi/6+2\pi(k/3)i}, \quad k = 0, 1, 2.$$

Equivalently, we may write

$$2^{1/3}e^{i\pi/6}, 2^{1/3}e^{5\pi i/6}, \text{ and } 2^{1/3}e^{9\pi i/6}.$$

These, too, are evenly spaced around a circle (although in this case the circle has radius $2^{1/3}$).

It would appear that the n^{th} roots of any complex number are always spaced evenly around a circle. Equivalently, their sum always seems to be zero. In fact, a more general result holds.

Theorem. Let $\xi_j := e^{2\pi ij/n}$, $j = 0, 1, 2, \dots, n-1$ be the n^{th} roots of unity.

Then

$$\frac{1}{n} \sum_{j=0}^{n-1} \xi_j^k = \delta_{k,0} = \begin{cases} 1, & \text{if } k = 0 \pmod{n}, \\ 0, & \text{otherwise.} \end{cases}$$

Proof. The case $k = 0$ is evident, so suppose $k \neq 0$. Recall the expression for the sum of the *finite geometric series*

$$\sum_{j=0}^{N-1} x^j = \frac{1-x^N}{1-x},$$

which can easily be proved by multiplying both sides by $(1-x)$. Then we

have

$$\sum_{j=0}^{n-1} \xi_j^k = \sum_{j=0}^{n-1} e^{2\pi i j k/n} = \sum_{j=0}^{n-1} (e^{2\pi i k/n})^j = \frac{1 - e^{2\pi i k}}{1 - e^{2\pi i k/n}} = 0,$$

because when $k \neq 0 \pmod n$ the numerator vanishes but denominator does not. □